

HEAT STRESS AND ATHLETIC PARTICIPATION

Early fall football, cross country, soccer and field hockey practices are conducted in very hot and humid weather in many parts of the United States. Due to the equipment and uniform needed in football, most of the heat problems have been associated with football. Heatstroke deaths in football were the highest in 1997 since the 1978 season. There is no excuse for heatstroke deaths to increase if the proper precautions are taken. Under such conditions, the athlete is subject to the following:

HEAT CRAMPS -- Painful cramps involving abdominal muscles and extremities caused by intense, prolonged exercise in the heat and depletion of salt and water due to profuse sweating.

HEAT SYNCOPE -- Weakness, fatigue and fainting due to loss of salt and water in sweat and exercise in the heat. Predisposes to heatstroke.

HEAT EXHAUSTION (WATER DEPLETION) -- Excessive weight loss, reduced sweating, elevated skin and deep body temperature, excessive thirst, weakness, headache and sometimes unconsciousness.

HEAT EXHAUSTION (SALT DEPLETION) -- Exhaustion, nausea, vomiting, muscle cramps, and dizziness due to profuse sweating and inadequate replacement of body salts.

HEATSTROKE -- An acute medical emergency related to thermoregulatory failure. Associated with nausea, seizures, disorientation, and possible unconsciousness or coma. It may occur suddenly without being preceded by any other clinical signs. The individual is usually unconscious with a high body temperature and a hot dry skin (heatstroke victims, contrary to popular belief, may sweat profusely). It is believed that the above mentioned heat stress problems can be controlled provided certain precautions are taken. According to the American Academy of Pediatrics Committee on Sports Medicine, heat related illnesses are all preventable. (Sports Medicine: Health Care for Young Athletes, American Academy of Pediatrics, 1991). The following practices and precautions are recommended:

1. Each athlete should have a physical exam with a medical history when first entering a program and an annual health history update. History of previous heat illness and type of training activities before organized practice begins should be included. State high school association's recommendations should be followed.
2. It is clear that top physical performance can only be achieved by an athlete who is in top physical condition. Lack of physical fitness impairs the performance of an athlete who participates in high temperatures. Coaches should know the **PHYSICAL CONDITION** of their athletes and set practice schedules accordingly.
3. Along with physical conditioning, the factor of acclimatization to heat is important. Acclimatization is the process of becoming adjusted to heat and it is essential to provide for **GRADUAL ACCLIMATIZATION TO HOT WEATHER**. It is necessary for an athlete to exercise in the heat if he/she is to become acclimatized to it. It is suggested that a graduated physical conditioning program be used and that 80% acclimatization can be expected to occur after the first 7 to 10 days. Final stages of acclimatization to heat are marked by increased sweating and reduced salt concentration in the sweat.
4. The old idea that water should be withheld from athletes during workouts has **NO SCIENTIFIC FOUNDATION**. The most important safeguard to the health of the athlete is the replacement of water. Water must be on the field and readily available to the athletes at all times. It is recommended that a minimum of 10 minutes be scheduled for a water break every half hour of heavy exercise in the heat. **WATER SHOULD BE AVAILABLE IN UNLIMITED QUANTITIES**. Check and be sure athletes are drinking the water. Cold water is preferable. Drinking ample water before practice or games has also been found to aid performance in the heat.
5. Salt should be replaced daily. Modest salting of foods after practice or games will accomplish this purpose. Salt tablets are not recommended. **ATTENTION MUST BE DIRECTED TO REPLACING WATER -- FLUID REPLACEMENT IS ESSENTIAL**.
6. Know both the **TEMPERATURE AND HUMIDITY**. The greater the humidity, the more difficult it is for the body to cool itself. Test the air prior to practice or game using a wet bulb, globe, temperature index (WBGT Index) which is based on the combined effects of air temperature, relative humidity, radiant heat and air movement. The following precautions are recommended when using the WBGT Index (ACSM's Guidelines for the Team Physician, 1991):

Below 64	Unlimited activity
65 - 72	Moderate risk
74 - 82	High risk
82 plus	Very high risk

There is also a weather guide for activities that last 30 minutes or more (Fox and Mathews, 1981) which involves knowing the relative humidity and air temperature:

<u>Air Temp</u>	<u>Danger Zone</u>	<u>Critical Zone</u>
70F	80% RH	100% RH
75F	70% RH	100% RH
80F	50% RH	80% RH
85F	40% RH	68% RH
90F	30% RH	55% RH
95F	20% RH	40% RH
100F	10% RH	30% RH

RH = Relative Humidity

One other method of measuring the relative humidity is the use of a sling psychrometer, which measures wet bulb temperature. The wet bulb temperature should be measured prior to practice and the intensity and duration of practice adjusted accordingly. Recommendations are as follows:

Under 60F Safe but always observe athletes
61 - 65F Observe players carefully
66 - 70F Caution
71 - 75F Shorter practice sessions and more frequent water and rest breaks
75 plus F Danger level and extreme caution

7. Cooling by evaporation is proportional to the area of the skin exposed. In extremely hot and humid weather reduce the amount of clothing covering the body as much as possible. **NEVER USE RUBBERIZED CLOTHING.**
8. Athletes should weight each day before and after practice and **WEIGHT CHARTS CHECKED.** Generally a 3 percent weight loss through sweating is safe and over a e percent weight loss is in the danger zone. Over a 3 percent weight loss the athlete should not be allowed to practice in hot and humid conditions. Observe the athletes closely under all conditions.
9. Observe athletes carefully for signs of trouble, particularly athletes who lose much weight and the eager athlete who constantly competes at his/her capacity. Some trouble signs are nausea, incoherence fatigue, weakness, vomiting, cramps, weak rapid pulse, visual disturbance and unsteadiness.
10. Teams that encounter hot weather during the season through travel or following an unseasonably cool period, should be physically fit, but will not be environmentally fit. Coaches in this situation should follow the above recommendations and substitute more frequently during games.
11. Know what to do in case of an emergency and have your emergency plans written with copies to all your staff. Be familiar with immediate first aid practice and prearranged procedures for obtaining medical care, including ambulance service.

HEAT STROKE -- THIS IS A MEDICAL EMERGENCY -- DELAY COULD BE FATAL. Immediately cool body while waiting for transfer to a hospital. Remove clothing and use cool water on body. An increasing number of medical personnel are now using a treatment for heat illness that involves applying either alcohol or cool water to the victim's skin and vigorously fanning the body. The fanning causes evaporation and cooling. (Source -- The First Aider -- September 1987)

HEAT EXHAUSTION -- OBTAIN MEDICAL CARE AT ONCE. Cool body as you would for heat stroke while waiting for transfer to hospital. Give fluids if athlete is able to swallow and is conscious.

SUMMARY -- The main problem associated with exercising in the hot weather is water loss through sweating. Water loss is best replaced by allowing the athlete unrestricted access to water. Water breaks two or three times per hour are better than one break an hour. Probably the best method is to have water available at all times and to allow the athlete to drink water whenever he/she needs it. Never restrict the amount of water an athlete drinks, and be sure the athletes are drinking the water. The small amount of salt lost in sweat is adequately replaced by salting food at meals. Talk to your medical personnel concerning emergency treatment plans.